

# Odd Graceful Labeling of the Revised Friendship Graphs

E. M. Badr

Department of scientific computing,  
 Faculty of Computers & Information,  
 Benha University, Benha, Egypt,

## ABSTRACT

The aim of this paper is to present some odd graceful graphs. In particular we show that the revised friendship graphs  $F(kC_4)$ ,  $F(kC_8)$ ,  $F(kC_{12})$ ,  $F(kC_{16})$  and  $F(kC_{20})$  are odd graceful where  $k$  is any positive integer. Finally, we introduce a new conjecture " The revised friendship graph  $F(kC_n)$  is odd graceful where  $k$  is any positive integer and  $n = 0 \pmod{4}$  ).

## Keywords

Graph Theory, odd graceful labeling, friendship graphs.

## 1. INTRODUCTION

A graph  $G$  of size  $q$  is odd-graceful, if there is an injection  $\phi$  from  $V(G)$  to  $\{0, 1, 2, \dots, 2q-1\}$  such that, when each edge  $xy$  is assigned the label or weight  $|\phi(x) - \phi(y)|$ , the resulting edge labels are  $\{1, 3, 5, \dots, 2q-1\}$ . This definition was introduced in 1991 by Gnanajothi [1] who proved that the class of odd graceful graphs lies between the class of graphs with  $\alpha$ -labelings and the class of bipartite graphs. Gnanajothi [1] proved that every cycle  $C_n$  is odd graceful if  $n$  is even. It is known that the graphs which contain odd cycles are not odd graceful so Badr [2] used the subdivision notation for odd cycle in order to prove that the subdivision of linear triangular snakes are odd graceful. Badr et al [3] proved that the subdivision of ladders  $S(L_n)$  is odd graceful.

Rosa [4] proved that the cycle  $C_n$  is graceful if and only if  $n \equiv 0$  or  $3 \pmod{4}$ . Solairaju and Muruganatham [5] proved that the revised friendship graphs  $F(kC_3)$ ,  $F(kC_5)$  and  $F(2kC_3)$  are all even vertex graceful, where  $k$  is any positive integer.

**Definition 1.1:** A revised friendship graph  $F(kC_n)$ ,  $n \geq 3$  is defined as a connected graph containing  $k$  copies of  $C_n$  with a vertex in common.

### Example 1.2:




Figure 1: the revised friendship graphs  $F(4C_4)$  &  $F(2C_8)$

In this paper we show that the revised friendship graphs  $F(kC_4)$ ,  $F(kC_8)$ ,  $F(kC_{12})$ ,  $F(kC_{16})$  and  $F(kC_{20})$  are odd graceful where  $k$  is any positive integer. Finally, we introduce a new conjecture " The revised friendship graph  $F(kC_n)$  is odd graceful where  $k$  is any positive integer and  $n = 0 \pmod{4}$  ).

## 2. THE MAIN RESULTS

**Theorem 2.1:** The revised friendship graph  $F(kC_4)$  is odd graceful, where  $k$  is any positive integer.

### Proof:

Let  $G = F(kC_4)$  has  $q$  edges and  $p$  vertices. The graph  $G$  consists of the vertices  $u, u_1u_2u_3\dots u_{2k}$  and  $v_1v_2v_3\dots v_k$ , where the graph  $G$  consisting  $k$  copies of  $C_4$  with a vertex  $u$  in common, such that  $u_i$  is put between  $u$  and  $v_j$  where  $i = 1, 2, 3, \dots, 2k$  and  $j = 1, 2, 3, \dots, k$ . The graph  $G$  has  $q = 4k$  and  $p = 3k + 1$ , as shown in Figure 2.




Figure 2: the revised friendship graph  $F(kC_4)$

Define  $\phi: V(G) \rightarrow \{0, 1, 2, \dots, 2q-1\}$  as following:

$$\phi(u) = 0$$

$$\phi(u_i) = 2q - 2i + 1, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(v_i) = 2q - 8i + 4, \quad i = 1, 2, 3, \dots, k$$

$$a) \text{Max}_{v \in V(G)} \phi(v) = \max\{0, \max_{1 \leq i \leq 2k} 2q - 2i + 1, \max_{1 \leq i \leq k} (2q - 8i + 4)\}$$

$$= 2q - 1, \text{ the maximum value of all odds}$$

Hence,  $\phi(v) \in \{0, 1, 2, \dots, 2q-1\}$

b) Clearly, The function  $\phi$  is one-to-one mapping from the vertex set of  $G$  to the set  $\{0, 1, 2, \dots, 2q-1\}$

c) It remains to show that the labels of the edges of  $G$  are all the odd integers of the interval  $[1, 2q-1]$  The range of  $|\phi(u_i) - \phi(u)| = \{2q - 2i + 1 : i = 1, 2, \dots, 2k\}$

$$= \{2q - 1, 2q - 3 \dots 2q - 4k + 1\}$$


The range of  $|\phi(v_i) - \phi(u_{2i-1})| = \{4i - 1 \quad : i = 1, 2 \dots k\}$   
 $= \{3, 7 \dots 4k - 1\}$

The range of  $|\phi(v_i) - \phi(u_{2i})| = \{4i - 3 \quad : i = 1, 2 \dots k\}$   
 $= \{1, 5 \dots 4k - 3\}$

Hence,  $\{|\phi(u) - \phi(v)| : u, v \in E(G)\} = \{1, 3, 5 \dots 2q - 1\}$ .

So the revised friendship graph  $F(kC_4)$  is odd graceful.

**Example 2.2**




**Figure 3: the odd graceful labeling of the revised friendship graph  $F(5C_4)$ .**

**Theorem 2.3:** The revised friendship graph  $F(kC_6)$  is odd graceful, where  $k$  is any positive integer.

**Proof:**

Let  $G = F(kC_6)$  has  $q$  edges and  $p$  vertices. The graph  $G$  consists of the vertices  $u, u_1u_2u_3 \dots u_{2k}, x_1x_2x_3 \dots x_{2k}$  and  $v_1v_2v_3 \dots v_k$ , where the graph  $G$  consisting  $k$  copies of  $C_6$  with a vertex  $u$  in common, such that the vertex  $u$  is the common vertex,  $x_i$  is put between  $u$  and  $v_j$ ,  $u_i$  is put between  $u$  and  $x_j$ , such that  $i = 1, 2, 3 \dots 2k, j = 1, 2, 3 \dots k$ . The graph  $G$  has  $q = 6k$  and  $p = 5k + 1$ , as shown in Figure 4.



**Figure 4: the revised friendship graph  $F(kC_6)$**

Define  $\phi : V(G) \rightarrow \{0, 1, 2 \dots 2q - 1\}$  as following:

$$\begin{aligned} \phi(u) &= 0 \\ \phi(u_i) &= 2q - 2i + 1, \quad i = 1, 2, 3 \dots 2k \\ \phi(x_i) &= (4/3)q - 4i + 2, \quad i = 1, 2, 3 \dots 2k \\ \phi(v_i) &= (2/3)q - 4i + 3, \quad i\text{-odd} (i = 1, 3, 5 \dots) \\ \phi(v_i) &= (2/3)q - 4i + 1, \quad i\text{-even} (i = 2, 4, 6 \dots) \end{aligned}$$

a)

$$\begin{aligned} \text{Max}_{v \in V(G)} \phi(v) &= \max\{0, \max_{1 \leq i \leq 2k} (2q - 2i + 1), \max_{1 \leq i \leq 2k} ((4/3)q - 4i + 2), \\ &\max_{i=1,3,\dots,k}^{i\text{-odd}} ((2/3)q - 4i + 3), \max_{i=2,4,\dots,k}^{i\text{-even}} ((2/3)q - 4i + 1)\} \end{aligned}$$

$= 2q - 1$ , the maximum value of all odds

Hence,  $\phi(v) \in \{0, 1, 2 \dots 2q - 1\}$

b) Clearly, The function  $\phi$  is one-to-one mapping from the vertex set of  $G$  to the set  $\{0, 1, 2 \dots 2q - 1\}$

c) It remains to show that the labels of the edges of  $G$  are all the odd integers of the interval  $[1, 2q - 1]$  and that's as following:

The range of  $|\phi(u_i) - \phi(u)| = \{2q - 2i + 1, i = 1, 2 \dots 2k\}$   
 $= \{2q - 1, 2q - 3 \dots 2q - 4k + 1\}$

The range of  $|\phi(u_i) - \phi(x_i)| = \{(2/3)q + 2i - 1, i = 1, 2 \dots 2k\}$   
 $= \{(2/3)q + 1, (2/3)q + 3 \dots (2/3)q + 4k - 1\}$

The range of  $|\phi(v_i) - \phi(x_{2i-1})| = \{(2/3)q - 4i + 3, i\text{-odd} (i = 1, 3, 5 \dots, k)\}$

$$= \{(2/3)q - 1, (2/3)q - 9 \dots\}$$

The range of  $|\phi(v_i) - \phi(x_{2i})| = \{(2/3)q - 4i - 1, i\text{-odd} (i = 1, 3, 5 \dots, k)\}$

$$= \{(2/3)q - 5, (2/3)q - 13 \dots\}$$

The range of  $|\phi(v_i) - \phi(x_{2i-1})| = \{(2/3)q - 4i + 5, i\text{-even} (i = 2, 4, 6 \dots, k - 1)\}$

$$= \{(2/3)q - 3, (2/3)q - 11 \dots\}$$

The range of  $|\phi(v_i) - \phi(x_{2i})| = \{(2/3)q - 4i + 1, i\text{-even} (i = 2, 4, 6 \dots, k - 1)\}$

$$= \{(2/3)q - 7, (2/3)q - 15 \dots\}$$

Hence,  $\{|\phi(u) - \phi(v)| : uv \in E(G)\} = \{1, 3, 5 \dots 2q - 1\}$ .

So the revised friendship graph  $F(kC_6)$  is odd graceful.

**Theorem 2.4:** The revised friendship graph  $F(kC_8)$  is odd graceful, where  $k$  is any positive integer.

**Proof:**

Let  $G = F(kC_8)$  has  $q$  edges and  $p$  vertices. The graph  $G$  consists of the vertices  $u, u_1u_2u_3 \dots u_{2k}, x_1x_2x_3 \dots x_{2k}, h_1h_2h_3 \dots h_{2k}$  and  $v_1v_2v_3 \dots v_k$ , where the graph  $G$  consisting  $k$  copies of  $C_8$  with a vertex  $u$  in common, such that the vertex  $u$  is the common vertex,  $h_i$  is put between  $u$  and  $v_j$ ,  $x_i$  is put between  $u$  and  $h_i$ ,  $u_i$  is put between  $u$  and  $x_j$  such that  $i = 1, 2, 3 \dots 2k$  and  $j = 1, 2, 3 \dots k$ . The graph  $G$  has  $q = 8k$  and  $p = 7k + 1$ , as shown in the next figure.




Figure 5: the revised friendship graph  $F(kC_8)$

Define  $\phi: V(G) \rightarrow \{0, 1, 2, \dots, 2q-1\}$  as following:

$$\begin{aligned} \phi(u) &= 0 \\ \phi(u_i) &= 2q - 2i + 1, \quad i = 1, 2, 3, \dots, 2k \\ \phi(x_i) &= q - 4i + 2, \quad i = 1, 2, 3, \dots, 2k \\ \phi(h_i) &= 2q - 2i - 4k + 1, \quad i = 1, 2, 3, \dots, 2k \\ \phi(v_i) &= 2q - 8i + 4, \quad i = 1, 2, 3, \dots, k \end{aligned}$$

a)

$$\begin{aligned} \text{Max}_{v \in V(G)} \phi(v) &= \max\{0, \max_{1 \leq i \leq 2k} (2q - 2i + 1), \max_{1 \leq i \leq 2k} (q - 4i + 2), \\ &\quad \max_{1 \leq i \leq 2k} (2q - 2i - 4k + 1), \max_{1 \leq i \leq k} (2q - 8i + 4)\} \\ &= 2q - 1, \text{ the maximum value of all odds} \end{aligned}$$

Hence,  $\phi(v) \in \{0, 1, 2, \dots, 2q-1\}$

b) Clearly, The function  $\phi$  is one-to-one mapping from the vertex set of  $G$  to the set  $\{0, 1, 2, \dots, 2q-1\}$

c) It remains to show that the labels of the edges of  $G$  are all the odd integers of the interval  $[1, 2q-1]$  and that's as following:

$$\begin{aligned} \text{The range of } |\phi(u_i) - \phi(u)| &= \{2q - 2i + 1, i = 1, 2, \dots, 2k\} \\ &= \{2q - 1, 2q - 3, \dots, 2q - 4k + 1\} \end{aligned}$$

$$\begin{aligned} \text{The range of } |\phi(u_i) - \phi(x_i)| &= \{q + 2i - 1, i = 1, 2, \dots, 2k\} \\ &= \{q + 1, q + 3, \dots, q + 4k - 1\} \end{aligned}$$

$$\begin{aligned} \text{The range of } |\phi(h_i) - \phi(x_i)| &= \{q + 2i - 4k - 1, i = 1, 2, \dots, 2k\} \\ &= \{q - 4k + 1, q - 4k + 3, \dots, q - 1\} \end{aligned}$$

$$\begin{aligned} \text{The range of } |\phi(v_i) - \phi(h_{2i-1})| &= \{4k - 4i + 1, i = 1, 2, 3, \dots, k\} \\ &= \{4k - 3, 4k - 7, \dots, 1\} \end{aligned}$$

$$\begin{aligned} \text{The range of } |\phi(v_i) - \phi(h_{2i})| &= \{4k - 4i + 3, i = 1, 2, 3, \dots, k\} \\ &= \{4k - 1, 4k - 5, \dots, 3\} \end{aligned}$$

Hence,  $\{|\phi(u) - \phi(v)| : uv \in E(G)\} = \{1, 3, 5, \dots, 2q-1\}$ .

So the revised friendship graph  $F(kC_8)$  is odd graceful.

**Theorem 2.5:** The revised friendship graph  $F(kC_{12})$  is odd graceful, where  $k$  is any positive integer.

**Proof:**

Let  $G = F(kC_{12})$  has  $q$  edges and  $p$  vertices. The graph  $G$  consists of the vertices  $u, u_1u_2u_3 \dots u_{2k}, x_1x_2x_3 \dots x_{2k}, h_1h_2h_3 \dots h_{2k}, a_1a_2a_3 \dots a_{2k}, b_1b_2b_3 \dots b_{2k}$  and  $v_1v_2v_3 \dots v_k$ , where the graph  $G$  consisting  $k$  copies of  $C_{12}$  with a vertex  $u$  in common, such that the vertex  $u$  is the common vertex,  $b_i$  is put between  $u$  and  $v_j$ ,  $a_i$  is put between  $u$  and  $b_i$ ,  $h_i$  is put between  $u$  and  $a_i$ ,  $x_i$  is put between  $u$  and  $h_i$ ,  $u_i$  is put between  $u$  and  $x_i$  where  $i = 1, 2, 3, \dots, 2k$  and  $j = 1, 2, 3, \dots, k$ . The graph  $G$  has  $q = 12k$  and  $p = 11k + 1$ , as shown in the next figure.




Figure 6: the revised friendship graph  $F(kC_{12})$

Define  $\phi: V(G) \rightarrow \{0, 1, 2, \dots, 2q-1\}$  as following:

$$\begin{aligned} \phi(u) &= 0 \\ \phi(u_i) &= 2q - 2i + 1, \quad i = 1, 2, 3, \dots, 2k \\ \phi(x_i) &= (4/6)q - 4i + 2, \quad i = 1, 2, 3, \dots, 2k \\ \phi(h_i) &= 2q - 2i - 4k + 1, \quad i = 1, 2, 3, \dots, 2k \\ \phi(a_i) &= (8/6)q - 4i + 2, \quad i = 1, 2, 3, \dots, 2k \\ \phi(b_i) &= (2/6)q - 2i + 1, \quad i = 1, 2, 3, \dots, 2k \\ \phi(v_i) &= (4/6)q - 8i + 4, \quad i = 1, 2, 3, \dots, k \end{aligned}$$

a)

$$\begin{aligned} \text{Max}_{v \in V(G)} \phi(v) &= \max\{0, \max_{1 \leq i \leq 2k} (2q - 2i + 1), \max_{1 \leq i \leq 2k} ((4/6)q - 4i + 2), \\ &\quad \max_{1 \leq i \leq 2k} (2q - 2i - 4k + 1), \max_{1 \leq i \leq 2k} ((8/6)q - 4i + 2), \\ &\quad \max_{1 \leq i \leq 2k} ((2/6)q - 2i + 1), \max_{1 \leq i \leq k} ((4/6)q - 8i + 4)\} \\ &= 2q - 1, \text{ the maximum value of all odds} \end{aligned}$$

Hence,  $\phi(v) \in \{0, 1, 2, \dots, 2q-1\}$

b) Clearly, The function  $\phi$  is one-to-one mapping from the vertex set of  $G$  to the set  $\{0, 1, 2, \dots, 2q-1\}$

c) It remains to show that the labels of the edges of  $G$  are all the odd integers of the interval  $[1, 2q-1]$  and that's as following:

$$\text{The range of } |\phi(u_i) - \phi(u)| = \{2q - 2i + 1, i = 1, 2, \dots, 2k\}$$

$$= \{2q - 1, 2q - 3 \dots 2q - 4k + 1\}$$

The range of  $|\phi(u_i) - \phi(x_i)| = \{(4/3)q + 2i - 1, i = 1, 2 \dots 2k\}$

$$= \{(4/3)q + 1, (4/3)q + 3 \dots (4/3)q + 4k - 1\}$$

The range of  $|\phi(h_i) - \phi(x_i)| = \{(4/3)q + 2i - 4k - 1, i = 1, 2 \dots 2k\}$

$$= \{(4/3)q - 4k + 1, (4/3)q - 4k + 3 \dots (4/3)q - 1\}$$

The range of  $|\phi(h_i) - \phi(a_i)| = \{(2/3)q + 2i - 4k - 1, i = 1, 2 \dots 2k\}$

$$= \{(2/3)q - 4k + 1, (2/3)q - 4k + 3, \dots (2/3)q - 1\}$$

The range of  $|\phi(a_i) - \phi(b_i)| = \{q - 2i + 1, i = 1, 2 \dots 2k\}$

$$= \{q - 1, q - 3 \dots q - 4k + 1\}$$

The range of  $|\phi(v_i) - \phi(b_{2i-1})| = \{(1/3)q - 4i + 1, i = 1, 2, 3 \dots k\}$

$$= \{(1/3)q - 3, (1/3)q - 7 \dots 1\}$$

The range of  $|\phi(v_i) - \phi(b_{2i})| = \{(1/3)q - 4i + 3, i = 1, 2, 3 \dots k\}$

$$= \{(1/3)q - 1, (1/3)q - 5 \dots 3\}$$


Hence,  $\{|\phi(u) - \phi(v)| : u, v \in E(G)\} = \{1, 3, 5 \dots 2q - 1\}$ .

So the revised friendship graph  $F(kC_{12})$  is odd graceful.

**Theorem 2.6:** The revised friendship graph  $F(kC_{16})$  is odd graceful, where  $k$  is any positive integer.

**Proof:**

Let  $G = F(kC_{16})$  has  $q$  edges and  $p$  vertices. The graph  $G$  consists of the vertices  $u, u_1u_2u_3 \dots u_{2k}, x_1x_2x_3 \dots x_{2k}, h_1h_2h_3 \dots h_{2k}, a_1a_2a_3 \dots a_{2k}, b_1b_2b_3 \dots b_{2k}, c_1c_2c_3 \dots c_{2k}, d_1d_2d_3 \dots d_{2k}$  and  $v_1v_2v_3 \dots v_k$ , where the graph  $G$  consisting  $k$  copies of  $C_{16}$  with a vertex  $u$  in common, such that the vertex  $u$  is the common vertex,  $d_i$  is put between  $u$  &  $v_j$ ,  $c_i$  is put between  $u$  and  $d_i$ ,  $b_i$  is put between  $u$  and  $c_i$ ,  $a_i$  is put between  $u$  and  $b_i$ ,  $h_i$  is put between  $u$  and  $a_i$ ,  $x_i$  is put between  $u$  and  $h_i$ ,  $u_i$  is put between  $u$  and  $x_i$  where  $i = 1, 2, 3 \dots 2k$  and  $j = 1, 2, 3 \dots k$ . The graph  $G$  has  $q = 16k$  and  $p = 15k + 1$ , as shown in the next figure.



**Figure 7: the revised friendship graph  $F(kC_{16})$**

Define  $\phi: V(G) \rightarrow \{0, 1, 2 \dots 2q-1\}$  as following:

$$\phi(u) = 0$$

$$\phi(u_i) = 2q - 2i + 1, \quad i = 1, 2, 3 \dots 2k$$

$$\phi(x_i) = (1/2)q - 4i + 2, \quad i = 1, 2, 3 \dots 2k$$

$$\phi(h_i) = 2q - 2i - 4k + 1, \quad i = 1, 2, 3 \dots 2k$$

$$\phi(a_i) = q - 4i + 2, \quad i = 1, 2, 3 \dots 2k$$

$$\phi(b_i) = (1/4)q - 2i + 1, \quad i = 1, 2, 3 \dots 2k$$

$$\phi(c_i) = (3/2)q - 4i + 2, \quad i = 1, 2, 3 \dots 2k$$

$$\phi(d_i) = q - 2i + 1, \quad i = 1, 2, 3 \dots 2k$$

$$\phi(v_i) = (5/4)q - 8i + 4, \quad i = 1, 2, 3 \dots k$$

a)

$$\text{Max } \phi(v) = \max\{0, \max_{1 \leq i \leq 2k} (2q - 2i + 1), \max_{1 \leq i \leq 2k} ((1/2)q - 4i + 2),$$

$$\max_{1 \leq i \leq 2k} (2q - 2i - 4k + 1), \max_{1 \leq i \leq 2k} (q - 4i + 2),$$

$$\max_{1 \leq i \leq 2k} ((1/4)q - 2i + 1), \max_{1 \leq i \leq 2k} ((3/2)q - 4i + 2),$$

$$\max_{1 \leq i \leq 2k} (q - 2i + 1), \max_{1 \leq i \leq k} ((5/4)q - 8i + 4)\}$$

$$= 2q - 1, \text{ the maximum value of all odds}$$

Hence,  $\phi(v) \in \{0, 1, 2 \dots 2q - 1\}$

b) Clearly, The function  $\phi$  is one-to-one mapping from the

vertex set of  $G$  to the set  $\{0, 1, 2 \dots 2q - 1\}$

c) It remains to show that the labels of the edges of  $G$  are all the odd integers of the interval  $[1, 2q-1]$  and that's as following:

The range of  $|\phi(u_i) - \phi(u)| = \{2q - 2i + 1, i = 1, 2 \dots 2k\}$

$$= \{2q - 1, 2q - 3 \dots 2q - 4k + 1\}$$

The range of  $|\phi(u_i) - \phi(x_i)| = \{(3/2)q + 2i - 1, i = 1, 2 \dots 2k\}$

$$= \{(3/2)q + 1, (3/2)q + 3 \dots (3/2)q + 4k - 1\}$$

The range of  $|\phi(h_i) - \phi(x_i)| = \{(3/2)q + 2i - 4k - 1, i = 1, 2 \dots 2k\}$

$$= \{(3/2)q - 4k + 1, (3/2)q - 4k + 3 \dots (3/2)q - 1\}$$

The range of  $|\phi(h_i) - \phi(a_i)| = \{q + 2i - 4k - 1, i = 1, 2 \dots 2k\}$

$$= \{q - 4k + 1, q - 4k + 3 \dots q - 1\}$$

The range of  $|\phi(a_i) - \phi(b_i)| = \{(3/4)q - 2i + 1, i = 1, 2 \dots 2k\}$

$$= \{(3/4)q - 1, (3/4)q - 3 \dots (3/4)q - 4k + 1\}$$

The range of  $|\phi(c_i) - \phi(b_i)| = \{(5/4)q - 2i + 1, i = 1, 2 \dots 2k\}$

$$= \{(5/4)q - 1, (5/4)q - 3 \dots (5/4)q - 4k + 1\}$$

The range of  $|\phi(c_i) - \phi(d_i)| = \{(1/2)q - 2i + 1, i = 1, 2 \dots 2k\}$

$$= \{(1/2)q - 1, (1/2)q - 3 \dots (1/2)q - 4k + 1\}$$

The range of  $|\phi(v_i) - \phi(d_{2i-1})| = \{(1/4)q - 4i + 1, i = 1, 2, 3 \dots k\}$

$$= \{(1/4)q - 3, (1/4)q - 7 \dots (1/4)q - 4k + 1\}$$

The range of  $|\phi(v_i) - \phi(d_{2i})| = \{(1/4)q - 4i + 3, i=1,2,3 \dots k\}$   
 $= \{(1/4)q - 1, (1/4)q - 5 \dots (1/4)q - 4k + 3\}$


Hence,  $\{|\phi(u) - \phi(v)| : u, v \in E(G)\} = \{1, 3, 5 \dots 2q - 1\}$ .

So the revised friendship graph  $F(kC_{16})$  is odd graceful.

**Theorem 2.7:** The revised friendship graph  $F(kC_{20})$  is odd graceful, where  $k$  is any positive integer.

**Proof:**

Let  $G = F(kC_{20})$  has  $q$  edges and  $p$  vertices. The graph  $G$  consists of the vertices  $u, u_1u_2u_3 \dots u_{2k}, x_1x_2x_3 \dots x_{2k}, h_1h_2h_3 \dots h_{2k}, a_1a_2a_3 \dots a_{2k}, b_1b_2b_3 \dots b_{2k}, c_1c_2c_3 \dots c_{2k}, d_1d_2d_3 \dots d_{2k}, z_1z_2z_3 \dots z_{2k}, g_1g_2g_3 \dots g_{2k}$  and  $v_1v_2v_3 \dots v_k$ , where the graph  $G$  consisting  $k$  copies of  $C_{20}$  with a vertex  $u$  in common, such that the vertex  $u$  is the common vertex,  $g_i$  is put between  $u$  and  $v_j$ ,  $z_i$  is put between  $u$  and  $g_i$ ,  $d_i$  is put between  $u$  and  $z_i$ ,  $c_i$  is put between  $u$  and  $d_i$ ,  $b_i$  is put between  $u$  and  $c_i$ ,  $a_i$  is put between  $u$  and  $b_i$ ,  $h_i$  is put between  $u$  and  $a_i$ ,  $x_i$  is put between  $u$  and  $h_i$ ,  $u_i$  is put between  $u$  and  $x_i$  where  $i = 1, 2, 3, \dots, 2k$  and  $j = 1, 2, 3, \dots, k$ . The graph  $G$  has  $q = 20k$  and  $p = 19k + 1$ , as shown in the next figure.



**Figure 8:** the revised friendship graph  $F(kC_{20})$

Define  $\phi: V(G) \rightarrow \{0, 1, 2, \dots, 2q-1\}$  as following:

$$\phi(u) = 0$$

$$\phi(u_i) = 2q - 2i + 1, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(x_i) = (2/5)q - 4i + 2, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(h_i) = 2q - 2i - 4k + 1, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(a_i) = (4/5)q - 4i + 2, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(b_i) = 2q - 2i - 8k + 1, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(c_i) = (6/5)q - 4i + 2, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(d_i) = (7/5)q - 2i + 1, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(z_i) = (8/5)q - 4i + 2, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(g_i) = (1/5)q - 2i + 1, \quad i = 1, 2, 3, \dots, 2k$$

$$\phi(v_i) = q - 8i + 4, \quad i = 1, 2, 3, \dots, k$$

a)

$$\text{Max } \phi(v) = \max_{v \in V(G)} \{0, \max_{1 \leq i \leq 2k} (2q - 2i + 1), \max_{1 \leq i \leq 2k} ((2/5)q - 4i + 2),$$

$$\max_{1 \leq i \leq 2k} (2q - 2i - 4k + 1), \max_{1 \leq i \leq 2k} ((4/5)q - 4i + 2),$$

$$\max_{1 \leq i \leq 2k} (2q - 2i - 8k + 1), \max_{1 \leq i \leq 2k} ((6/5)q - 4i + 2),$$

$$\max_{1 \leq i \leq 2k} ((7/5)q - 2i + 1), \max_{1 \leq i \leq 2k} ((8/5)q - 4i + 2)$$

$$\max_{1 \leq i \leq 2k} ((1/5)q - 2i + 1), \max_{1 \leq i \leq k} (q - 8i + 4) \}$$

$$= 2q - 1, \text{ the maximum value of all odds}$$

Hence,  $\phi(v) \in \{0, 1, 2, \dots, 2q - 1\}$

b) Clearly, The function  $\phi$  is one-to-one mapping from the vertex set of  $G$  to the set  $\{0, 1, 2, \dots, 2q - 1\}$

c) It remains to show that the labels of the edges of  $G$  are all the odd integers of the interval  $[1, 2q-1]$  and that's as following:

$$\text{The range of } |\phi(u_i) - \phi(u)| = \{2q - 2i + 1, i = 1, 2, \dots, 2k\}$$

$$= \{2q - 1, 2q - 3, \dots, 2q - 4k + 1\}$$

$$\text{The range of } |\phi(u_i) - \phi(x_i)| = \{(8/5)q + 2i - 1, i = 1, 2, \dots, 2k\}$$

$$= \{(8/5)q + 1, (8/5)q + 3, \dots, (8/5)q + 4k - 1\}$$

$$\text{The range of } |\phi(h_i) - \phi(x_i)| = \{(8/5)q + 2i - 4k - 1, i = 1, 2, \dots, 2k\}$$

$$= \{(8/5)q - 4k + 1, (8/5)q - 4k + 3, \dots, (8/5)q - 1\}$$

$$\text{The range of } |\phi(h_i) - \phi(a_i)| = \{(6/5)q + 2i - 4k - 1, i = 1, 2, \dots, 2k\}$$

$$= \{(6/5)q - 4k + 1, (6/5)q - 4k + 3, \dots, (6/5)q - 1\}$$

$$\text{The range of } |\phi(a_i) - \phi(b_i)| = \{(6/5)q + 2i - 8k - 1, i = 1, 2, \dots, 2k\}$$

$$= \{(6/5)q - 8k + 1, (6/5)q - 8k + 3, \dots, (6/5)q - 4k - 1\}$$

$$\text{The range of } |\phi(c_i) - \phi(b_i)| = \{(4/5)q + 2i - 8k - 1, i = 1, 2, \dots, 2k\}$$

$$= \{(4/5)q - 8k + 1, (4/5)q - 8k + 3, \dots, (4/5)q - 4k - 1\}$$

$$\text{The range of } |\phi(c_i) - \phi(d_i)| = \{(1/5)q + 2i - 1, i = 1, 2, \dots, 2k\}$$

$$= \{(1/5)q + 1, (1/5)q + 3, \dots, (1/5)q + 4k - 1\}$$

$$\text{The range of } |\phi(d_i) - \phi(z_i)| = \{(1/5)q - 2i + 1, i = 1, 2, \dots, 2k\}$$

$$= \{(1/5)q - 1, (1/5)q - 3, \dots, 1\}$$

$$\text{The range of } |\phi(z_i) - \phi(g_i)| = \{(7/5)q - 2i + 1, i = 1, 2, \dots, 2k\}$$

$$= \{(7/5)q - 1, (7/5)q - 3, \dots, (7/5)q - 4k + 1\}$$

$$\text{The range of } |\phi(v_i) - \phi(g_{2i-1})| = \{(4/5)q - 4i + 1, i = 1, 2, 3, \dots, k\}$$

$$= \{(4/5)q - 3, (4/5)q - 7, \dots, (4/5)q - 4k + 1\}$$

$$\text{The range of } |\phi(v_i) - \phi(g_{2i})| = \{(4/5)q - 4i + 3, i = 1, 2, 3, \dots, k\}$$

$$= \{(4/5)q - 1, (4/5)q - 5, \dots, (4/5)q - 4k + 3\}$$

Hence,  $\{|\phi(u) - \phi(v)| : u, v \in E(G)\} = \{1, 3, 5, \dots, 2q - 1\}$ .

So the revised friendship graph  $F(kC_{20})$  is odd graceful.

Now, we introduce a new conjecture and that's as shown.

**Conjecture 2.8:** The revised friendship graph  $F(kC_n)$  is odd graceful where  $k$  is any positive integer and  $n = 0 \pmod{4}$ ".

### 3. Conclusion

Graceful and odd gracefulness of a graph are two entirely different concepts. A graph may possess one or both of these or neither. In this paper we introduced the odd graceful labeling of the revised friendship graphs  $F(kC_4)$ ,  $F(kC_8)$ ,  $F(kC_{12})$ ,  $F(kC_{16})$  and  $F(kC_{20})$  where  $k$  is any positive integer. Finally, we introduced a new conjecture " The revised friendship graph  $F(kC_n)$  is odd graceful where  $k$  is any positive integer and  $n = 0 \pmod{4}$  ).

### 4. References

- [1] R.B. Gnanajothi, Topics in graph theory, Ph.D. thesis, Madurai Kamaraj University, India, 1991.
- [2] E. M. Badr, *On the Odd Gracefulness of Cyclic Snakes With Pendant Edges*, International journal on applications of graph theory in wireless ad hoc networks and sensor networks (GRAPH-HOC) Vol.4, No.4, December 2012.
- [3] E. M. Badr, M. I. Moussa & K. Kathiresan (2011): *Crown graphs and subdivision of ladders are odd graceful*, International Journal of Computer Mathematics, 88:17, 3570-3576.
- [4] A. Rosa, On certain valuation of the vertices of a graph, Theory of Graphs (International Symposium, Rome, July 1966), Gordon and Breach, New York and Dunod Paris (1967) 349-355.
- [5] A. Solairaju & P. Muruganatham, Even Vertex Gracefulness of Fan Graph, International Journal of Computer Applications, Volume 8-No.8, October 2010.